

My Canadian Pharmacy about Drugs For Better Sex

Impaired potency is divided into two main types - erectile dysfunction and premature ejaculation. In response to a sexual stimulus, the walls of the blood vessels supplying the penis with blood are relaxed. As a result, the penis, or rather, its cavernous bodies, are filled with blood. However, if for some reason there is a spasm, then an erection does not occur or it is too weak.

Many enzymes are involved in this process. First of all, it is phosphodiesterase, as well as nitric oxide. Also, men often have problems with the duration of erection or rapid ejaculation. As a rule, they are caused by another reason - the lack of neurotransmitters in the nervous system.

Medications for ED: How to Build Close-Knit Family



Types of drugs to increase potency

There are several types of drugs for impotence given by My Canadian Pharmacy. Here are their main groups:

- phosphodiesterase inhibitors;
- NO synthase stimulants;
- alpha-blockers;
- antispasmodics;
- dietary supplements based on vegetable or animal components.

Most drugs are taken orally, in pill form. However, there are forms of injections and even sprays. Needless to say, in most cases, tablet dosage forms are much more convenient, and injections cannot always be done immediately before sexual intercourse.

Phosphodiesterase inhibitors

The most famous group of drugs includes phosphodiesterase inhibitors. The list of main drugs in the group includes:

- Sildenafil;
- Tadalafil;
- Udenafil;
- Vardenafil.

Sildenafil is dispensed under the tradename - Viagra. The drug manufacturer is the American pharmaceutical company, Pfizer. The drug advantages were explored by chance because the original drug was produced to treat the treatment of low blood pressure (hypotension). The action of the drug is rapid - within a few tens of minutes. This effect duration may be individual for each man. In general, it lasts up to 4 hours.

Though today more effective drugs that increase potency have been developed, Sildenafil still deserves its popularity. In addition to the original drug (Viagra), there are a certain number of generics based on the same active substance. The main difference between brand and generic is the price.

Cialis is another ED drug the main component of which is Tadalafil. This drug was developed later than Viagra. The duration of its action exceeds the duration of the action of Viagra and is equal to 36 hours. The onset of the effect is also higher than that of preparations based on Sildenafil, and is usually 30 minutes, in rare cases - an hour or more.

Levitra main component is Vardenafil. It is produced by the British pharmaceutical company Glaxo Smith Kline. The effect of the drug appears about 20-25 minutes and persists for about 4 hours. The benefit is a minimum of contraindications. It affects the liver, kidneys and cardiovascular system to a lesser extent compared to Viagra. Also, the drug can be combined with alcohol in moderate doses and fatty foods. However, in this case, the effect of the drug may occur somewhat later than usual.

However, the drugs from this group have a lot of contraindications. They are contraindicated in the case of heart failure, kidney failure, hypertension, diabetes and some other chronic diseases. Taking the drugs is strictly contraindicated in the case of simultaneous use of nitroglycerin.

In addition, the drugs will not help in the event of a violated libido, that is, sexual desire. Drugs are intended only to enhance erection, and not to restore sexual desire. Studies show that the effectiveness of drugs in this group is quite high and equal to 70-92%. These are relatively safe drugs, although they should not be taken without the advice of a doctor, especially by people with chronic diseases.